

## 11 Metrizable

Next we try to construct metrics on compact spaces. We shall learn that a compact space is metrizable if and only if the corresponding commutative  $C^*$ -algebra is separable. Metrizable is equivalent to the existence of a countable family of continuous functions separating the points of the space. As a vague analogy to the manifolds, the reader may view such a countable family as a set of coordinate functions on the space.

**Theorem.** *If  $\mathcal{F} \subset C(X)$  is a countable family separating the points of a compact space  $(X, \tau)$  then  $X$  is metrizable.*

**Proof.** Let  $\mathcal{F} = \{f_n\}_{n=0}^\infty \subset C(X)$  separate the points of  $X$ . We can assume that  $\|f_n\| \leq 1$  for every  $n \in \mathbb{N}$ ; otherwise consider for instance functions  $x \mapsto f_n(x)/(1 + |f_n(x)|)$ . Let us define

$$d(x, y) := \sup_{n \in \mathbb{N}} 2^{-n} |f_n(x) - f_n(y)|$$

for every  $x, y \in X$ . Next we prove that  $d : X \times X \rightarrow [0, \infty[$  is a metric:  $d(x, y) = 0 \Leftrightarrow x = y$ , because  $\{f_n\}_{n=0}^\infty$  is a separating family. Clearly also  $d(x, y) = d(y, x)$  for every  $x, y \in X$ . Let  $x, y, z \in X$ . We have the triangle inequality:

$$\begin{aligned} d(x, z) &= \sup_{n \in \mathbb{N}} 2^{-n} |f_n(x) - f_n(z)| \\ &\leq \sup_{n \in \mathbb{N}} (2^{-n} |f_n(x) - f_n(y)| + 2^{-n} |f_n(y) - f_n(z)|) \\ &\leq \sup_{m \in \mathbb{N}} 2^{-m} |f_m(x) - f_m(y)| + \sup_{n \in \mathbb{N}} 2^{-n} |f_n(y) - f_n(z)| \\ &= d(x, y) + d(y, z). \end{aligned}$$

Hence  $d$  is a metric on  $X$ .

Finally, let us prove that the metric topology coincides with the original topology,  $\tau_d = \tau$ : Let  $x \in X$ ,  $\varepsilon > 0$ . Take  $N \in \mathbb{N}$  such that  $2^{-N} < \varepsilon$ . Define

$$U_n := f_n^{-1}(\mathbb{D}(f_n(x), \varepsilon)) \in \mathcal{V}_\tau(x), \quad U := \bigcap_{n=0}^N U_n \in \mathcal{V}_\tau(x).$$

If  $y \in U$  then

$$d(x, y) = \sup_{n \in \mathbb{N}} 2^{-n} |f_n(x) - f_n(y)| < \varepsilon.$$

Thus  $x \in U \subset B_d(x, \varepsilon) = \{y \in X \mid d(x, y) < \varepsilon\}$ . This proves that the original topology  $\tau$  is finer than the metric topology  $\tau_d$ , i.e.  $\tau_d \subset \tau$ . Combined with the facts that  $(X, \tau)$  is compact and  $(X, \tau_d)$  is Hausdorff, this implies that we must have  $\tau_d = \tau$   $\square$

**Corollary.** *Let  $X$  be a compact Hausdorff space. Then  $X$  is metrizable if and only if it has a countable basis.*

**Proof.** Suppose  $X$  is a compact space, metrizable with a metric  $d$ . Let  $r > 0$ . Then  $\mathcal{B}_r = \{B_d(x, r) \mid x \in X\}$  is an open cover of  $X$ , thus having a finite subcover  $\mathcal{B}'_r \subset \mathcal{B}_r$ . Then  $\mathcal{B} := \bigcup_{n=1}^{\infty} \mathcal{B}'_{1/n}$  is a countable basis for  $X$ .

Conversely, suppose  $X$  is a compact Hausdorff space with a countable basis  $\mathcal{B}$ . Then the family

$$\mathcal{C} := \{(B_1, B_2) \in \mathcal{B} \times \mathcal{B} \mid \overline{B_1} \subset B_2\}$$

is countable. For each  $(B_1, B_2) \in \mathcal{C}$  Urysohn's Lemma provides a function  $f_{B_1 B_2} \in C(X)$  satisfying

$$f_{B_1 B_2}(\overline{B_1}) = \{0\} \quad \text{and} \quad f_{B_1 B_2}(X \setminus B_2) = \{1\}.$$

Next we show that the countable family

$$\mathcal{F} = \{f_{B_1 B_2} : (B_1, B_2) \in \mathcal{C}\} \subset C(X)$$

separates the points of  $X$ : Take  $x, y \in X$ ,  $x \neq y$ . Then  $W := X \setminus \{y\} \in \mathcal{V}(x)$ . Since  $X$  is a compact Hausdorff space, there exists  $U \in \mathcal{V}(x)$  such that  $\overline{U} \subset W$ . Take  $B', B \in \mathcal{B}$  such that  $x \in B' \subset \overline{B'} \subset B \subset U$ . Then  $f_{B' B}(x) = 0 \neq 1 = f_{B' B}(y)$ . Thus  $X$  is metrizable  $\square$

**Conclusion.** *Let  $X$  be a compact Hausdorff space. Then  $X$  is metrizable if and only if  $C(X)$  is separable (i.e. contains a countable dense subset).*

**Proof.** Suppose  $X$  is a metrizable compact space. Let  $\mathcal{F} \subset C(X)$  be a countable family separating the points of  $X$  (as in the proof of the previous Corollary). Let  $\mathcal{G}$  be the set of finite products of functions  $f$  for which  $f \in \mathcal{F} \cup \mathcal{F}^* \cup \{\mathbb{I}\}$ ; the set  $\mathcal{G} = \{g_j\}_{j=0}^{\infty}$  is countable. The linear span  $\mathcal{A}$  of  $\mathcal{G}$  is the involutive algebra generated by  $\mathcal{F}$  (the smallest  $*$ -algebra containing  $\mathcal{F}$ ); due to the Stone–Weierstrass Theorem,  $\mathcal{A}$  is dense in  $C(X)$ . If  $S \subset \mathbb{C}$  is a countable dense set then

$$\left\{ \lambda_0 \mathbb{I} + \sum_{j=1}^n \lambda_j g_j \mid n \in \mathbb{Z}^+, (\lambda_j)_{j=0}^n \subset S \right\}$$

is a countable dense subset of  $\mathcal{A}$ , thereby dense in  $C(X)$ .

Conversely, assume that  $\mathcal{F} = \{f_n\}_{n=0}^\infty \subset C(X)$  is a dense subset. Take  $x, y \in X$ ,  $x \neq y$ . By Urysohn's Lemma there exists  $f \in C(X)$  such that  $f(x) = 0 \neq 1 = f(y)$ . Take  $f_n \in \mathcal{F}$  such that  $\|f - f_n\| < 1/2$ . Then

$$|f_n(x)| < 1/2 \quad \text{and} \quad |f_n(y)| > 1/2,$$

so that  $f_n(x) \neq f_n(y)$ ;  $\mathcal{F}$  separates the points of  $X$  □

**Exercise\*.** Prove that a topological space with a countable basis is separable. Prove that a metric space has a countable basis if and only if it is separable.

**Exercise.** There are non-metrizable separable compact Hausdorff spaces! Prove that  $X$  is such a space, where

$$X = \{f : [0, 1] \rightarrow [0, 1] \mid x \leq y \Rightarrow f(x) \leq f(y)\}$$

is endowed with a relative topology. Hint: Tihonov's Theorem.