

EXAMPLES OF UNIFORMLY QUASIREGULAR MAPPINGS

Kirsi Peltonen



TEKNILLINEN KORKEAKOULU
TEKNISKA HÖGSKOLAN
HELSINKI UNIVERSITY OF TECHNOLOGY
TECHNISCHE UNIVERSITÄT HELSINKI
UNIVERSITE DE TECHNOLOGIE D'HELSINKI

Kirsi Peltonen: Examples of uniformly quasiregular mappings; Helsinki University of Technology Institute of Mathematics Research Reports A417 (1999).

Abstract: *In this paper we construct examples of uniformly quasiregular (uqr) mappings. These provide counterexamples for a rigidity conjecture in quasiregular dynamics. It states that a closed manifold of dimension at least three, which admits a branched uqr mapping, is quasiconformally equivalent to an ordinary sphere S^n .*

AMS subject classifications: 30C65, 58F (primary), 53C (secondary).

Keywords: uniformly quasiregular mappings, quasiregular dynamics, branched covering.

ISBN 951-22-4637-6

ISSN 0784-3143

Painotalo Libella Oy, Espoo, 1999

Helsinki University of Technology
Department of Engineering Physics and Mathematics
Institute of Mathematics
P.O. Box 1100, 02015 HUT, Finland
email: *math@hut.fi*
downloadables: *http://www.math.hut.fi/*

author's email: *Kirsi.Peltonen@hut.fi*

1 Statements and results

In this paper we construct examples of a certain type of quasiregular mappings. Topologically, quasiregular mappings are branched coverings, that is, covering mappings without the local injectivity requirement. This is a consequence of the completely analytic definition demanding that the mapping distorts the metric only by a bounded amount. More precisely:

Definition 1 *A continuous mapping $f : G \rightarrow \mathbf{R}^n$ of a domain $G \subset \mathbf{R}^n$ is quasiregular, if*

$$f \text{ belongs to } W_{n,loc}^1(G)$$

and if there exists a number K , $1 \leq K < \infty$, such that

$$|f'(x)|^n \leq K J_f(x)$$

holds almost everywhere in G .

In the above definition $W_{n,loc}^1(G)$ is the Sobolev space of mappings with weak first order partial derivatives which are locally L^n integrable. For the above mappings the partial derivatives in the ordinary sense exist almost everywhere. We can thus define the formal derivative of f in terms of partial derivatives. The Jacobian determinant $\det f'(x)$ of f at x is denoted by $J_f(x)$. By the norm of the linear mapping $f'(x)$ we mean here the operator norm

$$|f'(x)| = \sup_{|h|=1} |f'(x)h|.$$

The above definition generalizes immediately to Riemannian manifolds. The branch set B_f is the set of points for which f is not locally homeomorphic. For the basic properties and theory of quasiregular mappings see [R].

Definition 2 *A mapping $g : M^n \rightarrow M^n$ is uniformly quasiregular if all its m -fold iterates g^m (compositions with itself) are quasiregular with a uniform bound on the dilatation.*

For the motivation of this terminology coming from groups of quasiconformal mappings and basic properties see [IM] and [M]. The theory of quasiregular dynamics is further developed in [MM].

Theorem 3 *Let M^n be a spherical space form. Then M^n admits a branched uniformly quasiregular mapping.*

A spherical space form M^n is a smooth complete, connected Riemannian manifold of constant sectional curvature $K > 0$. These can be characterized as spaces isometric to quotients S^n/Γ , where $S^n = S^n(\frac{1}{\sqrt{K}}) \subset \mathbf{R}^{n+1}$ is equipped with the standard metric and $\Gamma \subset O(n+1)$ acts freely and properly discontinuously [W]. If we consider M^n equipped with this metric giving positive constant curvature, the corresponding covering map $h : S^n \rightarrow M^n$

is an isometry whenever an injection. Any smooth manifold M^n with S^n as a universal covering space and fundamental group Γ acting properly discontinuously on S^n gives a spherical space form isometric to S^n/Γ . Complete classification of spherical space forms is given in [W].

It turns out that the metric is not an obstruction here and the situation is even more general. Namely we have the following.

Theorem 4 *Let M^n be a smooth Riemannian manifold with universal covering space S^n . Then M^n admits a branched uniformly quasiregular mapping.*

Theorem 5 *Let M^n be as in Theorem 4. Then any quasiregular mapping $f : M^n \rightarrow S^n$ composed by the covering mapping $S^n \rightarrow M^n$ can be deformed to a uniformly quasiregular mapping $M^n \rightarrow M^n$.*

In the above theorem by a deformation of a quasiregular mapping $M^n \rightarrow M^n$ we mean a uniformly quasiregular mapping $M^n \rightarrow M^n$ with the same branch set.

Theorem 6 *Theorems 4 and 5 hold if in place of a covering mapping there is a branched covering mapping.*

Corollary 7 *The Poincaré homology sphere and the Lens spaces in dimension 3 admit uniformly quasiregular mappings.*

The proof of the above statements is by construction and essentially a modification of a trick invented by G. Martin and T. Iwaniec [IM], [M]. We construct an analogous conformal trap, where a ball is conformally mapped into itself. The two essential steps in building the trap in [M] are the following:

- (1) A local modification of the original quasiregular mapping $S^n \rightarrow S^n$ to become a translation of the trap outside itself.
- (2) Inversion with respect to the boundary of the trap.

Step one generalizes immediately to a quasiregular map $M^n \rightarrow S^n$. The only global property needed in the construction is the inversion (2). This step we can take in a suitably chosen ball which is mapped further onto the trap by the covering mapping.

The dynamics of these mappings is the same as for maps in [M]. See also [HM].

Acknowledgements. This paper was born after an inspiring lecture by Gaven Martin in Haifa, May 1999, on open questions in the field of uqr mappings, and patient explanations by Volker Mayer on some related constructions. The author wants to thank the both gentlemen on dynamical discussions.

2 Proofs

Proof of Theorem 3 Assume M is a spherical space form of constant sectional curvature $K > 0$. Denote by $S^n = S^n(\frac{1}{\sqrt{K}}) \subset \mathbf{R}^{n+1}$ the sphere of radius $\frac{1}{\sqrt{K}}$ centered at the origin of \mathbf{R}^{n+1} with the standard metric. We equip M^n with the natural metric for which $h : S^n \rightarrow M^n = S^n/\Gamma$ is a local isometry. Then we take an Alexander type mapping $f : M^n \rightarrow S^n$, which is constructed as follows. Divide the compact manifold into finitely many simplices and further take a barycentric subdivision by adding a new vertex into each of the faces. In this situation there is always an even number of simplices attached to each of the $(n-2)$ -faces. Consider then the sphere S^n divided into two by upper and lower hemispheres. The simplices of the manifold M^n are then mapped pairwise to the two pieces of the sphere respectively. The branch set of the mapping is exactly the $(n-2)$ -skeleton. Assume that the degrees of the mappings f and h are p and q respectively. The construction works as well for any other quasiregular mapping available from M^n onto S^n in place of f .

Let $x_0 \in M^n$ be any point outside the branch set. Denote by y_0 one of the points in the set $h^{-1}(x_0)$. One can assume that $y_0 \neq f(x_0)$. We will build a conformal trap to the neighbourhood of the point x_0 . By making an inversion in the neighbourhood of the point y_0 analogously as in the construction in [M]. Analogously we need the following properties.

- (1) Set $f^{-1}(y_0)$ and x_0 do not meet B_f
- (2) There are small, disjoint balls $U_0 = B(f(x_0), r)$ and $V_0 = B(y_0, r)$ about $f(x_0)$ and y_0 in S^n such that $f^{-1}(U_0)$ has a component U_1 containing x_0 and $f^{-1}(V_0)$ has components V_1, \dots, V_p pairwise disjoint and such that $f|_{U_1} : U_1 \rightarrow U_0$, $f|_{V_i} : V_i \rightarrow V_0$ and $h|_{V_0} : V_0 \rightarrow h(V_0)$ are injective.

The above can be organized for almost all pairs of points (x_0, y_0) [R]. Denote by $\{y_1, \dots, y_p\} = f^{-1}(y_0)$. Let $a, b > 0$ be so small that $2b < a$ and the following conditions are satisfied:

$$B(x_0, a) \subset U_1, B(y_i, a) \subset V_i, i = 1, \dots, p \quad (1)$$

$$B(y_0, b) \subset \cap_{i=1}^p f(B(y_i, a)), B(f(x_0), b) \subset f(B(x_0, a)) \quad (2)$$

$$f(B(x_0, b)) \subset B(f(x_0), a), f(B(y_i, b)) \subset B(y_0, a), i = 1, \dots, p \quad (3)$$

Now define an intermediary map g_1 as follows. On

$$M^n \setminus \{\cup_{i=1}^p B(y_i, a) \cup B(x_0, a)\}$$

we set $g_1 = f$. For $1 \leq i \leq p$ we map $B(x_0, b)$ onto $B(f(x_0), b)$ and $B(y_i, b)$ onto $B(y_0, b)$ isometrically. On the annular regions $g_1|_{B(x_0, a) \setminus B(x_0, b)}$ and $g_1|_{B(y_i, a) \setminus B(y_i, b)}$ is obtained for each $i = 1, \dots, p$ by application of Sullivan's version of the Annulus Theorem for quasiconformal mappings ([TV]).

In the above $g_1|_{\partial B(x_0,a)} = f$, $g_1|_{\partial B(y_i,a)} = f$ and $g_1|_{\partial B(x_0,b)}$, $g_1|_{\partial B(y_i,b)}$ is the appropriate isometry. The conditions (1)-(3) imply the map $g_1 : M^n \rightarrow S^n$ is well defined and quasiregular. Additionally we have by construction $B_{g_1} = B_f$.

Next let $\Phi : S^n \rightarrow S^n$ be a conformal mapping which exchanges $B(y_0, b)$ with its complement and set

$$g = h \circ \Phi \circ g_1 : M^n \rightarrow M^n. \quad (4)$$

Clearly $B_g = B_f$. Next we show that g and all its iterates are uniformly quasiregular. Denote by $B = B(x_0, b)$ the conformal trap. Restriction map $g|_B$ is conformal and

$$g(B) = h(\Phi(B(f(x_0), b))) \subset h(B(y_0, b)) = B(h(y_0), b) = B$$

by (3) and conformality of h . It follows that $g^m|_B = g \circ \dots \circ g|_B$ is conformal for all $m \geq 1$. Assume next that $x \in M^n \setminus \{\cup_{i=1}^p B(y_i, b) \cup B(x_0, b)\}$. Then $g_1(x) \in S^n \setminus B(y_0, b)$ and hence $\Phi(g_1(x)) \in B(y_0, b)$ and finally $g(x) \in B$ since h is an isometry in $B(y_0, b)$. It follows that $g^m|_{M^n \setminus \{\cup_{i=1}^p B(y_i, b) \cup B(x_0, b)\}}$ is quasiregular for each m with a uniform bound on the dilatation by our first observation. Finally if $x \in B(y_i, b)$, $i = 1, \dots, p$, then g is a conformal isometry followed by a conformal Φ and h as above. Thus the iterates stay conformal at x until it passes into the complement of $\cup_{i=1}^p B(y_i, b) \cup B(x_0, b)$. Then it picks up some distortion before passing into B and the iterates again stay conformal. \square

Proof of Theorem 4 If the manifold M^n is equipped with an arbitrary Riemannian metric, it would be tempting to take the induced metric by the covering mapping on the sphere, making the covering $h : S^n \rightarrow M^n$ conformal. However, in this situation the inversion Φ need not be conformal. This was pointed out by V. Mayer. We equip the sphere with the standard spherical metric and modify the covering mapping, which is locally quasiconformal, as we modified the mapping f in the proof of Theorem 3. We proceed exactly as in the proof of Theorem 3 and use the same notation. The modification for h needs to be done in small neighbourhoods of y_0 and $h^{-1}(y_i)$, $i = 1, \dots, p$. We do the following additions. Denote by $\{z_i^1, \dots, z_i^q\} = h^{-1}(y_i)$, $i = 1, \dots, p$. We choose $r > 0$ here so small that in addition to the conditions in the proof of Theorem 3, also sets $\{W_i^1, \dots, W_i^q\} = h^{-1}B(y_i, r)$, $i = 1, \dots, p$, and V_0 are pairwise disjoint and $h|_{W_i^j} : W_i^j \rightarrow h(W_i^j)$, $h|_{V_0} : V_0 \rightarrow h(V_0)$ are injective for all $i = 1, \dots, p$, $j = 1, \dots, q$. Further we let $a, b > 0$ be so small that also the following conditions are satisfied:

$$B(z_i^j, a) \subset W_i^j, j = 1, \dots, q, i = 1, \dots, p \quad (5)$$

$$B(y_i, b) \subset \cap_{i=1}^q h(B(z_i^j, a)), i = 1, \dots, p, B(x_0, b) \subset h(B(y_0, a)) \quad (6)$$

$$h(B(z_i^j, b)) \subset B(y_i, a), j = 1, \dots, q, i = 1, \dots, p \quad (7)$$

$$h(B(y_0, b)) \subset B(x_0, a) \quad (8)$$

We define an intermediary map h_1 on $S^n \setminus \{\cup_{i,j=1}^{p,q} B(z_i^j, a) \cup B(y_0, a)\}$ by setting $h_1 = h$. For $j = 1, \dots, q$ we map $B(z_i^j, b)$ onto $B(y_i, b)$ and $B(y_0, b)$ onto $B(x_0, b)$ isometrically for each $i = 1, \dots, p$ and extend to annular regions as in Theorem 3. We claim now that the mapping $g = h_1 \circ \Phi \circ g_1$ is uqr. The conformal trap and points in $M^n \setminus \{\cup_{i=1}^p B(y_i, b) \cup B(x_0, b)\}$ behave as in the proof of Theorem 3. If

$$x \in g_1^{-1}(\Phi^{-1}(h_1^{-1}(B(y_i, b)))) \subset \cup_{i=1}^p B(y_i, b),$$

then $g(x)$ ends up to the set $\cup_{i=1}^p B(y_i, b)$ under conformal steps. If

$$x \in B(y_i, b) \setminus g_1^{-1}(\Phi^{-1}(h_1^{-1}(B(y_i, b)))),$$

then

$$\Phi(g_1(x)) \in S^n \setminus \{B(y_0, b) \cup_{i=1}^p h_1^{-1}(B(y_i, b))\}$$

and $g(x)$ ends up to $M^n \setminus \cup_{i=1}^p B(y_i, b)$ picking up some dilatation under h_1 . After that g_1 can pick up some dilatation again but then $g_1(g(x)) \in S^n \setminus B(y_0, b)$ and $g^2(x) \in B(x_0, b)$ ending to the trap. \square

Proof of Theorem 5 The statement follows immediately since our construction does not make any explicit use of the Alexander mapping $M^n \rightarrow S^n$. \square

Proof of Theorem 6 The similar construction for the branched covering mapping $h : S^n \rightarrow M^n$ is possible by choosing the relevant points outside the branch set of the corresponding mappings. It is enough to choose x_0 outside the set $B_f \cup hB_h \cup hfhB_h$ and such that $f(x_0)$ is outside B_h . The set of such points is dense in M^n . In this situation none of the points $y_0, h^{-1}(y_i), i = 1, \dots, p$, belongs to B_h and h can be modified to be an isometry in small neighbourhoods of those points. \square

Proof of Corollary 7 The statement is evident. The Poincaré homology sphere and Lens spaces are interesting 3-dimensional examples of manifolds whose universal covering space is the 3-sphere. The Poincaré homology sphere has fundamental group of order 120. For its several descriptions see [KS] and a detailed dodecahedral construction see [P]. It is an example of a homogeneous spherical space form. Lens space $L(p, q) = S^3/(\mathbf{Z}/p)$ has fundamental group of order p . For these and more examples in dimension 3 see [Mo] and also in higher dimensions see [W]. \square

References

- [HM] Hinkkanen, A. and Martin, G.J. *Attractors in quasiregular semigroups*. Proc. XVI Nevanlinna colloquim, Eds. I. Laine and O. Martio, de Gruyter, Berlin-New York, 1996, 135-141.
- [IM] Iwaniec, T. and Martin, G.J. *Quasiregular semigroups* Ann. Acad. Sci. Fenn. 21, 1996, 241-254.
- [KS] Kirby, R. and Scharlemann M. *Eight faces of Poincaré homology sphere*. Proc. of Geometric Topology Conference, Athens, 1979.
- [M] Martin, G.J. *Branch sets of uniformly quasiregular maps*. Conformal Geometry and Dynamics. 1, 24-27.
- [MM] Martin, G.J. and Mayer, V. *Rigidity in holomorphic and quasiregular dynamics*. Preprint.
- [Mo] Montesinos, J.M. *Classical tessellations and three-manifolds*. Springer-Verlag, 1985
- [P] Peltonen, K. *Quasiregularity on homology manifolds*, MLI preprint no 14, 95/96.
- [R] Rickman, S. *Quasiregular mappings*, Springer-Verlag, 1993.
- [TV] Tukia, P. and Väisälä, J. *Lipschitz and quasiconformal approximation and extension*. Ann. Acad. Sci. Fenn. 6,1981, 303-342.
- [W] Wolf, J.A. *Spaces of constant curvature*. McGraw-Hill, 1967.

(continued from the back cover)

- A412 Marko Huhtanen
Ideal GMRES can be bounded from below by three factors, Jan 1999
- A411 Juhani Pitkäranta
The first locking-free plane-elastic finite element: historia mathematica, Jan 1999
- A410 Kari Eloranta
Bounded Triangular and Kagomé Ice, Jan 1999
- A409 Jukka Tuomela and Teijo Arponen
On the numerical solution of involutive ordinary differential systems: Boundary value problems, Dec 1998
- A408 Ville Turunen
Commutator Characterization of Periodic Pseudodifferential Operators, Dec 1998
- A407 Jarmo Malinen
Discrete time Riccati equations and invariant subspaces of linear operators, Feb 1999
- A406 Jarmo Malinen
Riccati Equations for H^∞ Discrete Time Systems: Part II, Feb 1999
- A405 Jarmo Malinen
Riccati Equations for H^∞ Discrete Time Systems: Part I, Feb 1999
- A404 Jarmo Malinen
Toeplitz Preconditioning of Toeplitz Matrices an Operator Theoretic Approach, Feb 1999
- A403 Saara Hyvönen and Olavi Nevanlinna
Robust bounds for Krylov method, Nov 1998
- A402 Saara Hyvönen
Growth of resolvents of certain infinite matrices, Nov 1998
- A401 Jukka Tuomela
On the Numerical Solution of Involutive Ordinary Differential Systems: Boundary value problems, Dec 1998
- A400 Seppo Hiltunen
Implicit functions from locally convex spaces to Banach spaces, Jan 1999

HELSINKI UNIVERSITY OF TECHNOLOGY INSTITUTE OF MATHEMATICS
RESEARCH REPORTS

The list of reports is continued inside. Electronical versions of the reports are available at <http://www.math.hut.fi/reports/> .

- A417 Kirsi Peltonen
Examples of uniformly quasiregular mappings, Jul 1999
- A416 Saara Hyvönen
On the Iterative Solution of Nonnormal Problems, Jul 1999
- A415 Ville Turunen
A smooth operator valued symbol analysis, Aug 1999
- A414 Marko Huhtanen
A stratification of the set of normal matrices, May 1999
- A413 Clement Ph., Gripenberg G. and Londen S-O
Regularity properties of solutions of functional evaluation equations, May 1999

ISBN 951-22-4637-6

ISSN 0784-3143

Painotalo Libella Oy, Espoo, 1999